

$SL(2; \mathbb{R})$ -REPRESENTATIONS OF A BRIESKORN HOMOLOGY 3-SPHERE

TERUAKI KITANO AND YOSHIKAZU YAMAGUCHI

ABSTRACT. We classify $SL(2; \mathbb{C})$ -representations of a Brieskorn homology 3-sphere. We show any irreducible representation into $SL(2; \mathbb{C})$ is conjugate to that into either $SU(2)$ or $SL(2; \mathbb{R})$. We also give a construction of $SL(2; \mathbb{R})$ -representations for a Brieskorn homology 3-sphere from $PSL(2; \mathbb{R})$ -representations of the base orbifold fundamental group.

1. INTRODUCTION

In [7], the second author studied the asymptotic behavior of the Reidemeister torsion for Seifert fibered spaces with $SL(2; \mathbb{C})$ -representations which are not realized by $SU(2)$ -ones. We are interested in finding out what is the difference between irreducible $SL(2; \mathbb{C})$ -representations and $SU(2)$ -ones in the cases of Brieskorn homology 3-spheres. We will see that irreducible $SL(2; \mathbb{C})$ -representations consists of $SU(2)$ -ones and $SL(2; \mathbb{R})$ -ones, up to conjugate. The latter ingredients are constructed by homomorphisms from the fundamental group of the base orbifold into $PSL(2; \mathbb{R})$. We give a concrete correspondence between $SL(2; \mathbb{R})$ -representations of a Brieskorn homology 3-sphere and $PSL(2; \mathbb{R})$ -representations of the fundamental group of the base orbifold. This correspondence is derived by covering maps in a sequence of Seifert fibered spaces.

Let $\Sigma(a_1, a_2, a_3)$ be a Brieskorn homology 3-sphere which is defined by

$$\Sigma(a_1, a_2, a_3) = \{(z_1, z_2, z_3) \in \mathbb{C}^3 \mid z_1^{a_1} + z_2^{a_2} + z_3^{a_3} = 0, |z_1|^2 + |z_2|^2 + |z_3|^2 = 1\}.$$

Here a_1, a_2, a_3 are pairwise coprime positive integers. We may assume a_2, a_3 are odd without the loss of generality.

Our main result in this paper is the following.

Theorem 1.1. *Let ρ be an irreducible representation of $\pi_1 \Sigma(a_1, a_2, a_3)$ into $SL(2; \mathbb{C})$. If ρ is not conjugate to an $SU(2)$ -representation, then ρ is conjugate to an $SL(2; \mathbb{R})$ -representation which is induced by a homomorphism from the fundamental group of the base orbifold into $PSL(2; \mathbb{R})$.*

2010 Mathematics Subject Classification. 57M25.

Key words and phrases. Brieskorn homology sphere, $SL(2; \mathbb{C})$ -representation, $SU(2)$ -representation, $PSL(2; \mathbb{R})$ -representation.

It is well known that $\Sigma(a_1, a_2, a_3)$ admits a structure of Seifert fibered space

$$\Sigma(a_1, a_2, a_3) \rightarrow S^2(a_1, a_2, a_3)$$

such that the base orbifold is $S^2(a_1, a_2, a_3)$, which is topologically S^2 with three cone points. We write $\Gamma(a_1, a_2, a_3)$ to the orbifold fundamental group of $S^2(a_1, a_2, a_3)$ with a presentation

$$\Gamma(a_1, a_2, a_3) = \langle x, y, z \mid x^{a_1} = y^{a_2} = z^{a_3} = xyz = 1 \rangle.$$

In this article, we consider irreducible representations of $\pi_1 \Sigma(a_1, a_2, a_3)$ into Lie groups $SL(2; \mathbb{C})$, $SU(2)$, $SL(2; \mathbb{R})$. For a Lie group G , we denote the space of conjugacy classes of irreducible representations of $\pi_1 \Sigma(a_1, a_2, a_3)$ into G by $\mathcal{R}_G$.

Remark 1.2. We consider conjugacy classes of irreducible representations under conjugation by $SL(2; \mathbb{C})$.

There are many studies on $\mathcal{R}_{SU(2)}$, and $\mathcal{R}_{SL(2; \mathbb{C})}$ which are related with $SU(2)$, or $SL(2; \mathbb{C})$ -Casson invariant of a homology 3-sphere.

In [3] Fintushel and Stern proved that $\mathcal{R}_{SU(2)}$ is a finite set. In particular the number is expressed as

$$|\mathcal{R}_{SU(2)}| = \frac{(a_1 - 1)(a_2 - 1)(a_3 - 1)}{4} - |X_0|$$

where the finite set X_0 is defined by

$$X_0 = \{(k, l, m) \mid 0 < k < a_1, 0 < l < a_2, 0 < m < a_3, \frac{k}{a_1} + \frac{l}{a_2} + \frac{m}{a_3} < 1\}.$$

They also proved that

$$|\mathcal{R}_{SU(2)}| = 2|\lambda(\Sigma(a_1, a_2, a_3))|$$

where $\lambda(\Sigma(a_1, a_2, a_3))$ is the Casson invariant of $\Sigma(a_1, a_2, a_3)$.

Remark 1.3. In fact they studied the space of conjugacy classes of $SU(2)$ -representations under conjugation by $SU(2)$ itself. However it is equal to $\mathcal{R}_{SU(2)}$ for a Brieskorn homology 3-sphere.

Curtis [2] defined the $SL(2; \mathbb{C})$ -Casson invariant for homology 3-spheres by counting conjugacy classes of irreducible $SL(2; \mathbb{C})$ -representations. Further Boden and Curtis [1] proved the equality that

$$\begin{aligned} \frac{(a_1 - 1)(a_2 - 1)(a_3 - 1)}{4} &= |\mathcal{R}_{SL(2; \mathbb{C})}| \\ &= \lambda_{SL(2; \mathbb{C})}(\Sigma(a_1, a_2, a_3)) \end{aligned}$$

by looking into representations of $\Gamma(2a_1, a_2, a_3)$ in $SL(2; \mathbb{C})$.

Hence Fintushel-Stern's formula can be expressed as

$$\lambda_{SL(2; \mathbb{C})}(\Sigma(a_1, a_2, a_3)) - 2|\lambda(\Sigma(a_1, a_2, a_3))| = |X_0|.$$

On the other hand, Jankins and Neumann [4] studied the space of homomorphisms from $\Gamma(a_1, a_2, a_3)$ into $PSL(2; \mathbb{R})$. It is not a finite set, but connected components can be classified by euler classes for central extensions of $\Gamma(a_1, a_2, a_3)$ by $\mathbb{Z}$, which belong to $H^2(\Gamma(a_1, a_2, a_3); \mathbb{Z})$. The number of connected components coincides with $2|X_0|$.

Corollary 1.4. *We have that*

$$\lambda_{SL(2; \mathbb{C})}(\Sigma(a_1, a_2, a_3)) - 2|\lambda(\Sigma(a_1, a_2, a_3))| = |\mathcal{R}_{SL(2; \mathbb{R})}|.$$

2. PRELIMINARIES

2.1. Seifert invariants. We say that a Seifert fibered space with three singular fibers over the sphere has a Seifert invariant

$$\{0; (1, b), (a_1, b_1), (a_2, b_2), (a_3, b_3)\}$$

where b and b_i are integers. Here 0 is the genus of $S^2(a_1, a_2, a_3)$. We write $M(0; (1, b), (a_1, b_1), (a_2, b_2), (a_3, b_3))$ to the Seifert fibered space by the above invariant. Please see [5, 6] for general theory of Seifert fibered spaces.

By using these invariants, the fundamental group $\pi_1 \Sigma(a_1, a_2, a_3)$ has the following presentation:

$$\langle x, y, z, h \mid h \text{ central}, x^{a_1} = h^{-b_1}, y^{a_2} = h^{-b_2}, z^{a_3} = h^{-b_3}, xyz = h^b \rangle.$$

We use the following convention of the euler number for a Seifert fibration.

Definition 2.1. For $M = M(0; (1, b), (a_1, b_1), (a_2, b_2), (a_3, b_3))$, we will denote the euler number by $e(M)$ defined as

$$e(M) = -\left(b + \frac{b_1}{a_1} + \frac{b_2}{a_2} + \frac{b_3}{a_3}\right) \in \mathbb{Q}.$$

Proposition 2.2. *For any $M = M(0; (1, b); (a_1, b_1), (a_2, b_2), (a_3, b_3))$, one has*

$$H_1(M; \mathbb{Z}) \simeq \mathbb{Z}/a|e(M)|$$

where $a = a_1 a_2 a_3$.

For the proof, see [5].

Corollary 2.3. *If $a|e(M)| = 1$, then M is a homology 3-sphere. In particular $M \cong \Sigma(a_1, a_2, a_3)$.*

Here and subsequently, we take $b, b_1, b_2, b_3 \in \mathbb{Z}$ as

$$a \left(b + \frac{b_1}{a_1} + \frac{b_2}{a_2} + \frac{b_3}{a_3} \right) = 1.$$

Since h is corresponding to the regular fiber of $\Sigma(a_1, a_2, a_3)$, one has a short exact sequence

$$1 \rightarrow \langle h \rangle \rightarrow \pi_1 \Sigma(a_1, a_2, a_3) \rightarrow \Gamma(a_1, a_2, a_3) = \pi_1 \Sigma(a_1, a_2, a_3) / \langle h \rangle \rightarrow 1.$$

Further this is a central extension of $\Gamma(a_1, a_2, a_3)$ by $\langle h \rangle \cong \mathbb{Z}$.

If one choose integers β and β_i such that

$$a \left| \beta + \frac{\beta_1}{a_1} + \frac{\beta_2}{a_2} + \frac{\beta_3}{a_3} \right| \neq 1,$$

then the corresponding Seifert fibered space

$$\overline{M} = M(0; (1, \beta), (a_1, \beta_1), (a_2, \beta_2), (a_3, \beta_3))$$

is not a homology 3-sphere. We use the symbols $\overline{M}$, β and β_i for a Seifert rational homology 3-sphere.

2.2. $SL(2; \mathbb{R})$ -representation. The projective group $PSL(2; \mathbb{R})$ is the quotient $SL(2; \mathbb{R}) / \{\pm I\}$ where I is the identity matrix. $PSL(2; \mathbb{R})$ acts on $\mathbb{RP}^1 \cong S^1$ as projectively automorphisms. Let $\widetilde{PSL}(2; \mathbb{R})$ be the universal cover of $PSL(2; \mathbb{R})$. $\widetilde{PSL}(2; \mathbb{R})$ also acts on $\mathbb{R}^1$ which is the universal cover of $\mathbb{RP}^1$. Then $\widetilde{PSL}(2; \mathbb{R})$ can be considered as a subgroup in the group of homeomorphisms of $\mathbb{R}$. For any $r \in \mathbb{R}$, we define

$$\text{sh}(r) : \mathbb{R} \ni x \mapsto x + r \in \mathbb{R}.$$

The center of $\widetilde{PSL}(2; \mathbb{R})$ is given by $\{\text{sh}(n) : \mathbb{R} \rightarrow \mathbb{R} \mid n \in \mathbb{Z}\} = \langle \text{sh}(1) \rangle \cong \mathbb{Z}$.

Lemma 2.4. *This short exact sequence*

$$0 \rightarrow \mathbb{Z} \rightarrow \widetilde{PSL}(2; \mathbb{R}) \rightarrow PSL(2; \mathbb{R}) \rightarrow 0$$

is a central extension of $PSL(2; \mathbb{R})$ by $\mathbb{Z} = \langle \text{sh}(1) \rangle$.

Recall that $\Gamma(a_1, a_2, a_3)$ is the quotient group of $\pi_1 \Sigma(a_1, a_2, a_3)$ by the center $\langle h \rangle$ with the following presentation:

$$\Gamma(a_1, a_2, a_3) = \langle x, y, z \mid x^{a_1} = y^{a_2} = z^{a_3} = xyz = 1 \rangle.$$

Given a homomorphism $f : \Gamma(a_1, a_2, a_3) \rightarrow PSL(2; \mathbb{R})$, we take the pull-back of the central extension

$$0 \rightarrow \mathbb{Z} \rightarrow \widetilde{PSL}(2; \mathbb{R}) \rightarrow PSL(2; \mathbb{R}) \rightarrow 0.$$

This gives a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z} & \longrightarrow & \widetilde{\Gamma}(a_1, a_2, a_3) & \longrightarrow & \Gamma(a_1, a_2, a_3) \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow f \\ 0 & \longrightarrow & \mathbb{Z} & \longrightarrow & \widetilde{PSL}(2; \mathbb{R}) & \longrightarrow & PSL(2; \mathbb{R}) \longrightarrow 0. \end{array}$$

Here $\widetilde{\Gamma}(a_1, a_2, a_3)$ has the following presentation:

$$\langle x, y, z, h \mid h \text{ central}, x^{a_1} = h^{-\beta_1}, y^{a_2} = h^{-\beta_2}, z^{a_3} = h^{-\beta_3}, xyz = h^\beta \rangle$$

The classification of the central extension of $\Gamma(a_1, a_2, a_3)$ is given by the euler class $eu(f)$ in

$$H^2(\Gamma(a_1, a_2, a_3); \mathbb{Z}) \cong \langle x_0, x_1, x_2, x_3 \mid a_1 x_1 = x_0, a_2 x_2 = x_0, a_3 x_3 = x_0 \rangle_{\mathbb{Z}}$$

such that $eu(f) = \beta x_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3$. By the relations, one can normalizes any element $\gamma \in H^2(\Gamma(a_1, a_2, a_3); \mathbb{Z})$ as

$$\gamma = \beta x_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3$$

such that $0 < \beta_1 < a_1, 0 < \beta_2 < a_2$ and $0 < \beta_3 < a_3$.

Applying [4, Theorem 1] to our case that a_1, a_2, a_3 are pairwise coprime, we have the following theorem.

Theorem 2.5 (Jankins-Neumann [4]). *For any $\gamma = \beta x_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3$, it equals $eu(f)$ for some $f : \Gamma(a_1, a_2, a_3) \rightarrow PSL(2; \mathbb{R})$ if and only if the following is true;*

- (a) $\beta = -1, \frac{\beta_1}{a_1} + \frac{\beta_2}{a_2} + \frac{\beta_3}{a_3} < 1$,
- either
- (b) $\beta = -2, \frac{\beta_1}{a_1} + \frac{\beta_2}{a_2} + \frac{\beta_3}{a_3} > 2$.

Now assume that such an $f : \Gamma(a_1, a_2, a_3) \rightarrow PSL(2; \mathbb{R})$ exists. Then there exists a central extension

$$0 \rightarrow \langle h \rangle \rightarrow \widetilde{\Gamma}(a_1, a_2, a_3) \rightarrow \Gamma(a_1, a_2, a_3) \rightarrow 1$$

defined by the pull-back of

$$0 \rightarrow \mathbb{Z} \rightarrow \widetilde{PSL}(2; \mathbb{R}) \rightarrow PSL(2; \mathbb{R}) \rightarrow 1$$

such that $\widetilde{\Gamma}(a_1, a_2, a_3) \ni h \mapsto \text{sh}(1) \in \widetilde{PSL}(2; \mathbb{R})$.

The following holds from the proof of Theorem 2.5 in [4]

Proposition 2.6. $\widetilde{\Gamma}(a_1, a_2, a_3)$ is the fundamental group of the Seifert fibered space with Seifert invariant $\{0; (1, \beta), (a_1, \beta_1), (a_2, \beta_2), (a_3, \beta_3)\}$.

Hence by composing $\tilde{\Gamma}(a_1, a_2, a_3) \rightarrow \widetilde{PSL}(2; \mathbb{R})$ with the projection from $\widetilde{PSL}(2; \mathbb{R})$ onto $SL(2; \mathbb{R})$, one has an $SL(2; \mathbb{R})$ -representation

$$\bar{\rho} : \pi_1 \bar{M} = \tilde{\Gamma}(a_1, a_2, a_3) \rightarrow SL(2; \mathbb{R}).$$

Lemma 2.7. *The $SL(2; \mathbb{R})$ -representation $\bar{\rho}$ is irreducible as an $SL(2; \mathbb{C})$ -representation.*

Proof. Suppose that $\bar{\rho}$ were a reducible $SL(2; \mathbb{C})$ -representation. We can move the image of $\pi_1 \bar{M}$ into the set of upper triangular matrices by conjugation. Under the homomorphism $\pi_1 \bar{M} \rightarrow \widetilde{PSL}(2; \mathbb{R})$, we see that x^{a_1} , y^{a_2} and z^{a_3} are sent to $\text{sh}(-\beta_1)$, $\text{sh}(-\beta_2)$ and $\text{sh}(-\beta_3)$ in $\widetilde{PSL}(2; \mathbb{R})$ respectively. The images of x , y and z are conjugate to $\text{sh}(-\beta_1/a_1)$, $\text{sh}(-\beta_2/a_2)$ and $\text{sh}(-\beta_3/a_3)$ in $\widetilde{PSL}(2; \mathbb{R})$. The projection from $\widetilde{PSL}(2; \mathbb{R})$ onto $SL(2; \mathbb{R})$ sends $\text{sh}(r)$ to a matrix of trace $2 \cos r\pi$. We can see that $\text{tr} \bar{\rho}(xyz) \neq \pm 2$ from that $\bar{\rho}(x)$, $\bar{\rho}(y)$, $\bar{\rho}(z)$ are upper triangular and that a_1, a_2, a_3 are pairwise coprime, $0 < \beta_i < a_i$ and $\sum_{i=1}^3 \beta_i/a_i < 1$. This is a contradiction to that $\bar{\rho}(xyz) = \bar{\rho}(h)^\beta = (-I)^\beta$. $\square$

Note that for $\bar{M} = M(0; (1, \beta), (a_1, \beta_1), (a_2, \beta_2), (a_3, \beta_3))$, it holds that

$$H_1(\bar{M}; \mathbb{Z}) \cong \mathbb{Z}/a|e(\bar{M})|$$

where $e(\bar{M})$ is the euler number of the Seifert fibration.

Since Brieskorn homology 3-sphere $\Sigma(a_1, a_2, a_3)$ is a Seifert fibered space $M(0; (1, b), (a_1, b_1), (a_2, b_2), (a_3, b_3))$, we can regard $\Sigma(a_1, a_2, a_3)$ as a universal abelian cover for each $\bar{M}$ with $a|e(\bar{M})| \neq 1$. We denote the induced homomorphism from $\pi_1 \Sigma(a_1, a_2, a_3)$ to $\pi_1 \bar{M}$ by p . If an euler class eu satisfies a condition in Theorem 2.5, then we have an $SL(2; \mathbb{R})$ -representation $\bar{\rho}$ of $\pi_1 \bar{M}$. We also have an $SL(2; \mathbb{R})$ -representation of $\pi_1 \Sigma(a_1, a_2, a_3)$ given by the pull-back $p^* \bar{\rho}$.

Hereafter we focus on $SL(2; \mathbb{R})$ -representations $p^* \bar{\rho}$ which are induced from euler classes eu satisfying that $\beta = -1$ and $\beta_1/a_1 + \beta_2/a_2 + \beta_3/a_3 < 1$.

Lemma 2.8. *We have a one-to-one correspondence from euler classes satisfying the condition (a) to those satisfying (b) in Theorem 2.5.*

Proof. This is given by an orientation reversing homeomorphism between Seifert fibered spaces. Let $\bar{M}$ be the Seifert fibered space corresponding to an euler class satisfying the condition (a). By changing the Seifert invariant, we can express the Seifert invariant of $-\bar{M}$ as

$$\begin{aligned} -\bar{M} &= -M(0; (1, -1), (a_1, \beta_1), (a_2, \beta_2), (a_3, \beta_3)) \\ &= M(0; (1, 1), (a_1, -\beta_1), (a_2, -\beta_2), (a_3, -\beta_3)) \\ &= M(0; (1, -2), (a_1, a_1 - \beta_1), (a_2, a_2 - \beta_2), (a_3, a_3 - \beta_3)). \end{aligned}$$

This Seifert invariant satisfies the condition (b). $\square$

We have an $SL(2; \mathbb{R})$ -representation of $\pi_1 \Sigma(a_1, a_2, a_3)$ by the pull-back induced from the universal abelian covering between Seifert fibered spaces. We write ρ_{eu} for the pull-back of an $SL(2; \mathbb{R})$ -representation of $\pi_1 \overline{M}$ corresponding to the euler class eu . We denote by Φ the induced correspondence between euler classes eu and the conjugacy classes of ρ_{eu} , i.e.,

$$\Phi: E \rightarrow \mathcal{R}_{SL(2; \mathbb{R})} (\subset \mathcal{R}_{SL(2; \mathbb{C})})$$

where E is the set of euler classes $eu \in H^2(\Gamma(a_1, a_2, a_3); \mathbb{Z})$ given by $eu = \beta x_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3$, $\beta = -1$, $0 < \beta_i < a_i$ ($i = 1, 2, 3$) and $\sum_{i=1}^3 (\beta_i / a_i) < 1$.

Remark 2.9. It is obvious that the set E can be identified with X_0 .

3. MAIN RESULT

In this section we prove the main theorem. Here M denotes a Brieskorn homology 3-sphere $\Sigma(a_1, a_2, a_3)$.

Theorem 3.1. *Let ρ be an irreducible $SL(2; \mathbb{C})$ -representation of $\pi_1 M$. If ρ is not conjugate to an $SU(2)$ -representation, then there exists a unique euler class $eu \in E$ such that the induced $SL(2; \mathbb{R})$ -representation ρ_{eu} is conjugate to ρ .*

Proof. It is shown in [1] that

$$|\mathcal{R}_{SL(2; \mathbb{C})}| = \frac{(a_1 - 1)(a_2 - 1)(a_3 - 1)}{4}.$$

Together with the result [3], we can see that $|\mathcal{R}_{SL(2; \mathbb{C})}| - |\mathcal{R}_{SU(2)}| = |E|$. It remains to prove that

- (1) any irreducible $SU(2)$ -representation can not be moved to $SL(2; \mathbb{R})$ -one under conjugation by a matrix in $SL(2; \mathbb{C})$,
- (2) the map Φ is one-to-one.

We divide these proofs into Proposition 3.5 and 3.12. □

Corollary 3.2. *We have that $|\mathcal{R}_{SL(2; \mathbb{C})}| = |\mathcal{R}_{SU(2)}| + |\mathcal{R}_{SL(2; \mathbb{R})}|$ by counting the conjugacy classes as $SL(2; \mathbb{C})$ -representations.*

Remark 3.3. It remains the same to count the conjugacy classes of irreducible $SU(2)$ -representations under the both conjugations by $SU(2)$ and $SL(2; \mathbb{C})$. In the case of $SL(2; \mathbb{R})$, it is not same.

Remark 3.4. There exists an example in which has no irreducible $SL(2; \mathbb{R})$ -representations. It holds for $\Sigma(2, 3, 5)$ that $|\mathcal{R}_{SL(2; \mathbb{C})}| = |\mathcal{R}_{SU(2)}|$.

The following proposition is folklore.

Proposition 3.5. *Let $\rho : \pi_1 \Sigma(a_1, a_2, a_3) \rightarrow SU(2)$ be an irreducible representation. If ρ is conjugate to an $SL(2; \mathbb{R})$ -representation by a matrix in $SL(2; \mathbb{C})$, then ρ is abelian.*

Proposition 3.5 follows from the next lemma.

Lemma 3.6. *Let $A, B \in SU(2)$. If there exists $P \in SL(2; \mathbb{C})$ such that $PAP^{-1}, PBP^{-1} \in SL(2; \mathbb{R})$, then $AB = BA$.*

Proof. We prove that if $P^{-1}AP$ and $P^{-1}BP$ are matrices in $SL(2; \mathbb{R})$, then $AB = BA$. It means that A and B are contained in the same maximal abelian subgroup in $SU(2)$. It is known that every matrix in $SU(2)$ can be diagonalizable by a matrix in $SU(2)$. We can assume A is a diagonal matrix. Suppose that $P^{-1}AP, P^{-1}BP \in SL(2; \mathbb{R})$. We write T_A to the linear fraction transform by A and so on. It is known that $T_{P^{-1}AP}$ has the fixed points given by a and $\bar{a}$ (the complex conjugate of a). The fixed points of T_A are 0 and ∞ . Since T_P sends a and $\bar{a}$ to 0 and ∞ respectively, we can express T_P as

$$T_P = e^{i\theta} \frac{z - a}{z - \bar{a}} \quad (\theta \in \mathbb{R}).$$

It is known that this T_P maps the real line to the unit circle in the complex plane. Together with that the real linear fractional transform $T_{P^{-1}BP}$ maps the real line onto itself, we can see that $T_B = T_P \circ T_{P^{-1}BP} \circ T_P^{-1}$ maps the unit circle onto itself. Hence T_B is expressed as

$$T_B = e^{i\varphi} \frac{z - \zeta}{-\bar{\zeta}z + 1} \quad (\varphi \in \mathbb{R}).$$

On the other hand, we can express T_B as

$$T_B = \frac{\xi z + \eta}{-\bar{\eta}z + \bar{\xi}}, \quad |\xi|^2 + |\eta|^2 = 1$$

since B is an element in $SU(2)$. Since η and ζ must be zero, B is also diagonal. This means that A and B are contained in the same maximal abelian subgroup in $SU(2)$. $\square$

Remark 3.7. We have shown a more general situation. Let F be a finitely generated group and $\rho : F \rightarrow SU(2)$ a representation. If ρ is conjugate to an $SL(2; \mathbb{R})$ -representation by a matrix in $SL(2; \mathbb{C})$, then ρ is abelian.

Under an irreducible representation $\rho : \pi_1 \Sigma(a_1, a_2, a_3) \rightarrow SL(2; \mathbb{C})$, we write the capital letter X for $\rho(x)$. Any conjugacy class in $\mathcal{R}_{SL(2; \mathbb{C})}$ can be parametrized by the triple of traces $(\text{tr } X, \text{tr } Y, \text{tr } XY)$. This triple is determined under a choice of integers b and b_i in the Seifert invariant.

Remark 3.8. These integers b, b_1, b_2, b_3 are not unique for given a_1, a_2, a_3 . As in [1], we can choose $b = 0$. We will require this assumption in the proof of our main result.

We may assume b_2 is even because a_2 is odd. If b_2 is odd, then we can replace b_1 and b_2 with $b_1 + a_1$ and $b_2 - a_2$ respectively. We will show that

Φ is a one-to-one correspondence between the euler class and the triple of traces. First we assume that integers b and b_i in the Seifert invariant of $\Sigma(a_1, a_2, a_3)$ satisfy that $b = 0$ and b_2, b_3 are even. Under this assumption, the presentation of $\pi_1 \Sigma(a_1, a_2, a_3)$ turns into

$$\langle x, y, z, h \mid h \text{ central}, x^{a_1} = h^{-b_1}, y^{a_2} = h^{-b_2}, z^{a_3} = h^{-b_3}, xyz = 1 \rangle.$$

The trace $\text{tr } Z$ equals to $\text{tr}(XY)^{-1} = \text{tr } XY$. The triple $(\text{tr } X, \text{tr } Y, \text{tr } XY)$ coincides with $(\text{tr } X, \text{tr } Y, \text{tr } Z)$.

The following two lemmas give a necessary condition for $(\text{tr } X, \text{tr } Y, \text{tr } Z)$ to coincide with $(\text{tr } X', \text{tr } Y', \text{tr } Z')$. We set integers q_i and r_i as

$$-a|e(\overline{M})|b_i = a_i q_i + r_i \quad (0 < r_i < a_i).$$

Lemma 3.9. *The trace of X under ρ_{eu} is expressed as*

$$\text{tr } X = (-1)^{q_1} 2 \cos \frac{\beta_1}{a_1} \pi$$

Similarly, the traces of Y and Z are also expressed as

$$\text{tr } Y = (-1)^{q_2} 2 \cos \frac{\beta_2}{a_2} \pi, \quad \text{tr } Z = (-1)^{q_3} 2 \cos \frac{\beta_3}{a_3} \pi.$$

Proof. By definition, the $SL(2; \mathbb{R})$ -representation $\rho_{eu} = p^* \overline{\rho}$ factors through $\pi_1 \Sigma(a_1, a_2, a_3) \rightarrow \pi_1 \overline{M} \rightarrow \widetilde{PSL}(2; \mathbb{R})$. We denote the images in $\widetilde{PSL}(2; \mathbb{R})$ of the generators $x, y, z \in \pi_1 \Sigma(a_1, a_2, a_3)$ by $\tilde{x}, \tilde{y}, \tilde{z}$ respectively. Since the order $|H(\overline{M}; \mathbb{Z})|$ is $a|e(\overline{M})|$, the covering degree between $\Sigma(a_1, a_2, a_3)$ and $\overline{M}$ also equals to $a|e(\overline{M})|$. The regular fiber $h \in \pi_1 \Sigma(a_1, a_2, a_3)$ is sent to $\text{sh}(a|e(\overline{M})|)$ in $\widetilde{PSL}(2; \mathbb{R})$. By the relation $x^{a_1} = h^{-b_1}$, we have the relation $\tilde{x}^{a_1} = \text{sh}(-a|e(\overline{M})|b_1)$ in $\widetilde{PSL}(2; \mathbb{R})$. Hence $\tilde{x}$ is conjugate to $\text{sh}(-a|e(\overline{M})|b_1/a_1)$.

Since the image of $\text{sh}(r)$ has trace $2 \cos r\pi$ under the projection onto $SL(2; \mathbb{R})$, we have that

$$(*) \quad \text{tr } X = 2 \cos \frac{-a|e(\overline{M})|b_1}{a_1} \pi = (-1)^{q_1} 2 \cos \frac{r_1}{a_1} \pi$$

where $-a|e(\overline{M})|b_1 = a_1 q_1 + r_1$ ($0 < r_1 < a_1$).

Finally we recall the relation that $a(b + \sum_{i=1}^3 b_i/a_i) = 1$. This relation implies that $a_2 a_3 b_1 \equiv 1 \pmod{a_1}$. Therefore we have that $-a|e(\overline{M})|b_1 \equiv \beta_1 a_2 a_3 b_1 \equiv \beta_1 \pmod{a_1}$. Together with (*), we obtain our claim. $\square$

Lemma 3.10. *If ρ_{eu} is conjugate to $\rho_{eu'}$, then $\beta_i = \beta'_i$ for all i or there is a unique i such that $\beta'_i = a_i - \beta_i$ and $\beta_j = \beta'_j$ for $j \neq i$.*

Proof. By Lemma 3.9, the equality $\text{tr } X = \text{tr } X'$, $\text{tr } Y = \text{tr } Y'$ and $\text{tr } Z = \text{tr } Z'$ give the constraints that $\beta_i = \beta'_i$ or $\beta'_i = a_i - \beta_i$ for each i . From another

constraints that $\beta_1/a_1 + \beta_2/a_2 + \beta_3/a_3 < 1$ and $\beta'_1/a_1 + \beta'_2/a_2 + \beta'_3/a_3 < 1$, we must have only one different β'_i from β_i if there exists. $\square$

Remark 3.11. The conclusions in Lemma 3.10 are not exclusive. When a_1 equals to $2\beta_1$, the case of $i = 1$ in the latter conclusion coincides with the former one.

Proposition 3.12. *The map $\Phi : E \rightarrow \mathcal{R}_{SL(2;\mathbb{R})}$ is one-to-one.*

Proof. We prove that if the triple $(\text{tr } X, \text{tr } Y, \text{tr } Z)$ equals to $(\text{tr } X', \text{tr } Y', \text{tr } Z')$, then $e = e'$. From Lemma 3.10 and Remark 3.11, it remains to show that if there exists a unique i such that $\beta'_i = a_i - \beta_i$, then $a_i = 2\beta_i$. Consequently, e coincides with e' . We will show that $a_1 = 2\beta_1$ under our assumption that $b = 0$, a_2, a_3 are odd and b_2, b_3 are even. Suppose that $\beta'_1 = \beta_1 - a_1$. By the equation (*), we can express $\text{tr } X$ as

$$\begin{aligned} \text{tr } X &= 2 \cos \frac{-a|e(\overline{M})|b_1}{a_1} \pi \\ &= 2 \cos \left(-b_1 a_2 a_3 + \frac{\beta_1 b_1 a_2 a_3}{a_1} + b_1 \beta_2 a_3 + b_1 a_2 \beta_3 \right) \pi \\ &= -2 \cos \left(\frac{\beta_1 b_1 a_2 a_3}{a_1} + b_1 \beta_2 a_3 + b_1 a_2 \beta_3 \right) \pi, \end{aligned}$$

which is due to that $b_1 a_2 a_3$ is odd. On the other hands, we can also express $\text{tr } X'$ as

$$\begin{aligned} \text{tr } X' &= 2 \cos \left(-\frac{\beta_1 b_1 a_2 a_3}{a_1} + b_1 \beta_2 a_3 + b_1 a_2 \beta_3 \right) \pi \\ &= 2 \cos \left(-\frac{\beta_1 b_1 a_2 a_3}{a_1} - b_1 \beta_2 a_3 - b_1 a_2 \beta_3 \right) \pi. \end{aligned}$$

Since we assume that $\text{tr } X = \text{tr } X'$, we have $\text{tr } X = 0$ which shows that $a_1 = 2\beta_1$ from Lemma 3.9. The same conclusion $a_1 = 2\beta_1$ can be drawn for the cases that $\beta'_i = a_i - \beta_i$ for $i = 2, 3$ by similar arguments. These cases are excluded by the constraints that $\sum_i \beta_i/a_i < 1$ and $\sum_i \beta'_i/a_i < 1$. $\square$

We can use the set of euler classes satisfying the condition (b) in Theorem 2.5. This is due to that the Seifert fibered space $-\overline{M}$ also induces the same triple of traces as that for $\overline{M}$. We can express an isomorphism $\varphi : \pi_1 \overline{M} \rightarrow \pi_1(-\overline{M})$ as

$$\varphi : h \mapsto h'^{-1}, x \mapsto x'h', y \mapsto y'h', z \mapsto z'h'.$$

Let ρ and ρ' be the induced $SL(2; \mathbb{R})$ -representations by $\overline{M}$ and $-\overline{M}$ respectively. The triple for ρ is equal to that of ρ' . Note that $e(-\overline{M}) = -e(\overline{M})$ and

h is sent to $\text{sh}(-a|e(-\overline{M})|)$ in $\widetilde{PSL}(2; \mathbb{R})$ under ρ' . By a similar argument in the proof of Lemma 3.9, we can show that

$$\text{tr } \rho'(x) = 2 \cos \frac{a|e(-\overline{M})|b_1}{a_1} \pi = (-1)^{-q_1} 2 \cos \left(\frac{-\beta_1}{a_1} \pi \right) = \text{tr } \rho(x).$$

Also we have $\text{tr } \rho'(y) = \text{tr } \rho(y)$ and $\text{tr } \rho'(z) = \text{tr } \rho(z)$.

4. EXAMPLES

4.1. The Brieskorn homology 3-sphere $\Sigma(2, 3, 6n + 1)$. We consider the $SL(2; \mathbb{R})$ -representations of $\pi_1 \Sigma(2, 3, 6n + 1)$. It is known that the Brieskorn homology 3-sphere $\Sigma(2, 3, 6n + 1)$ is obtained by $1/n$ -surgery along the torus knot of type $(2, 3)$ in S^3 . Here we assume that $n > 0$. The Casson invariant $|\lambda(\Sigma(2, 3, 6n + 1))|$ equals to n and the $SL(2; \mathbb{C})$ -Casson invariant $\lambda_{SL(2; \mathbb{C})}(\Sigma(2, 3, 6n + 1))$ equals to $3n$ by [1, Theorem 2.3]. Since the difference $\lambda_{SL(2; \mathbb{C})}(\Sigma(2, 3, 6n + 1)) - 2|\lambda(\Sigma(2, 3, 6n + 1))|$ is n , we have n conjugacy classes of irreducible $SL(2; \mathbb{R})$ -representations.

The euler classes of $\Gamma(2, 3, 6n + 1)$. There exist n euler classes $\beta x_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3 \in H^2(\Gamma(2, 3, 6n + 1))$ such that $\beta = -1$ and $\beta_1/2 + \beta_2/3 + \beta_3/(6n + 1) < 1$. They are given by

$$-x_0 + x_1 + x_2 + kx_3$$

where $1 \leq k \leq n$.

We denote the corresponding Seifert fibered space by $\overline{M}_k$. The order of $H_1(\overline{M}_k; \mathbb{Z})$ is given by $6(n - k) + 1$. Note that $\overline{M}_n$ is the Brieskorn homology 3-sphere $\Sigma(2, 3, 6n + 1)$. We can change the Seifert invariant of $\overline{M}_n$ from $\{0; (1, -1), (2, 1), (3, 1), (6n + 1, n)\}$ to $\{0; (1, 0), (2, 1), (3, -2), (6n + 1, n)\}$. The fundamental group of $\pi_1 \overline{M}_n$ is expressed as

$$\pi_1 \overline{M}_n = \langle x, y, z, h \mid h \text{ central}, x^2 = h^{-1}, y^3 = h^2, z^{6n+1} = h^{-n}, xyz = 1 \rangle$$

The triples of $\text{tr } X$, $\text{tr } Y$ and $\text{tr } Z$. By Lemma 3.9, we have the $SL(2; \mathbb{R})$ -representations induced from the euler classes. The corresponding triples

of traces are given by

$$\begin{aligned}
\operatorname{tr} X &= 2 \cos \left(-\frac{6(n-k)+1}{2} \pi \right) = 0, \\
\operatorname{tr} Y &= 2 \cos \left(\frac{2(6(n-k)+1)}{3} \pi \right) = -1, \\
\operatorname{tr} Z &= 2 \cos \left(-\frac{(6(n-k)+1)n}{6n+1} \pi \right) \\
&= 2 \cos \left(-\left(n-k + \frac{k}{6n+1} \right) \pi \right) \\
&= \begin{cases} 2 \cos \left(\frac{k}{6n+1} \pi \right) & n-k \text{ is even;} \\ 2 \cos \left(\pi - \frac{k}{6n+1} \pi \right) & n-k \text{ is odd.} \end{cases}
\end{aligned}$$

Note that the trace $\operatorname{tr} Z = 2 \cos \frac{\ell}{6n+1} \pi$ for an $SU(2)$ -representation must satisfy that $\ell \equiv n \pmod{2}$ and

$$\frac{1}{6} < \frac{\ell}{6n+1} < \frac{5}{6}.$$

We can express explicit $\operatorname{tr} Z$ for $SU(2)$ -representations as

$$\operatorname{tr} Z = 2 \cos \left(\frac{n+2}{6n+1} \pi \right), 2 \cos \left(\frac{n+4}{6n+1} \pi \right), \dots, 2 \cos \left(\frac{5n}{6n+1} \pi \right).$$

We have seen that $\Sigma(2, 3, 6n+1)$ appears in $\overline{M}_k$. There exist examples in which $\Sigma(a_1, a_2, a_3)$ does not appear in the sequence of Seifert fibered spaces induced by euler classes in E .

4.2. The Brieskorn homology 3-sphere $\Sigma(3, 5, 7)$. Under the constraint that $b = -1$ and $0 < b_1 < 3$, $0 < b_2 < 5$ and $0 < b_3 < 7$, we can choose the Seifert invariant as $\{0; (1, -1), (3, 2), (5, 1), (7, 1)\}$. Since $2/3 + 1/5 + 1/7 > 1$, $\Sigma(3, 5, 7)$ does appear in the sequence of $\overline{M}$. In fact, we have 4 euler classes satisfying that the condition (a) in Theorem 2.5. They are given by

$$\begin{aligned}
&-x_0 + x_1 + x_2 + x_3, & -x_0 + x_1 + x_2 + 2x_3, \\
&-x_0 + x_1 + x_2 + 3x_3, & -x_0 + x_1 + 2x_2 + x_3.
\end{aligned}$$

We can see that every corresponding Seifert fibered space $\overline{M}$ has the non-trivial homology group $H_1(\overline{M}; \mathbb{Z})$ by calculating the orders.

Acknowledgments. This paper was written while the authors were visiting Aix-Marseille University. They would like to express their sincere thanks for the hospitality. They also wish to express their thanks to Joan Porti for his helpful comments. This research was supported in part by JSPS KAKENHI 25400101 and 26800030.

REFERENCES

1. H. Boden and C. Curtis, *The $SL_2(\mathbb{C})$ Casson invariant for Seifert fibered homology spheres and surgeries on twist knots*, J. Knot Theory Ramifications **15** (2006), no. 7, 813–837.
2. C. Curtis, *An intersection theory count of the $SL_2(\mathbb{C})$ -representations of the fundamental group of a 3-manifold*, Topology **40** (2001), no. 4, 773–787. Erratum, Topology **42** (2003), no. 4, 929.
3. R. Fintushel and R. J. Stern, *Instanton homology of Seifert fibred homology three spheres*, Proc. London Math. Soc. (3) **61** (1990), no. 1, 109–137.
4. M. Jenkins and W. Neumann, *Homomorphisms of Fuchsian groups to $PSL(2, \mathbb{R})$* , Comment. Math. Helv. **60** (1985), no. 3, 480–495.
5. M. Jenkins and W. Neumann, *Lectures on Seifert manifolds*, Brandeis Lecture Notes, 2. Brandeis University, Waltham, MA, 1983. i+84+27 pp.
6. P. Orlik, *Seifert manifolds*, Lecture Notes in Mathematics, Vol. **291**. Springer-Verlag, Berlin-New York, 1972.
7. Y. Yamaguchi, *The asymptotic behavior of the Reidemeister torsion for Seifert manifolds and $PSL(2; \mathbb{R})$ -representations of Fuchsian groups*, arXiv:1410.2970

DEPARTMENT OF INFORMATION SYSTEMS SCIENCE, FACULTY OF SCIENCE AND ENGINEERING,
SOKA UNIVERSITY, TANGI-CHO 1-236, HACHIOJI, TOKYO 192-8577, JAPAN

E-mail address: kitano@soka.ac.jp

DEPARTMENT OF MATHEMATICS, AKITA UNIVERSITY, 1-1 TEGATA-GAKUENMACHI, AKITA, 010-
8502, JAPAN

E-mail address: shouji@math.akita-u.ac.jp